VOL. 593, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY 28 FEBRUARY, 1992

# CONTENTS

## 15TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, BASEL, JUNE 3–7, 1991, PART II

### COLUMN LIQUID CHROMATOGRAPHY (*Continued*)

*Applications: pharmacological*

Assessment of the biotransformation of the cardiotonic agent piroximone by high-performance liquid chromatography and gas chromatography–mass spectrometry
by M. Berg-Candolfi, J. T. Borlakoglu, B. Dulery, F. Jehl and K. D. Haegele (Strasbourg, France) . . . . . 1

Determination of amphetamines by high-performance liquid chromatography with ultraviolet detection. On-line pre-column derivatization with 9-fluorenylmethyl chloroformate and preconcentration
by G. Maeder, M. Pelletier and W. Haerdi (Geneva, Switzerland) . . . . . 9

High-performance liquid chromatographic determination of penicillin G, penicillin V and cloxacillin in beef and pork tissues
by W. A. Moats (Beltsville, MD, USA) . . . . . 15

Automated high-performance liquid chromatographic method for the determination of rifampicin in plasma
by K. J. Swart and M. Papgis (Bloemfontein, South Africa) . . . . . 21

High-performance liquid chromatographic analysis of FCE 24304 (6-methylenandrosta-1,4-diene-3,17-dione) and FCE 24928 (4-aminoandrosta-1,4,6-triene-3,17-dione), two new aromatase inhibitors
by S. Del Nero, M. Di Somma and A. Vigevani (Milan, Italy) . . . . . 25

High-performance liquid chromatographic determination of monohydroxy compounds by a combination of pre-column derivatization and post-column reaction detection
by V. K. Boppana, R. C. Simpson, K. Anderson, C. Miller-Stein, T. J. A. Blake, B. Y.-H. Hwang and G. R. Rhodes (King of Prussia, PA, USA) . . . . . 29

Direct high-performance liquid chromatographic resolution of a novel benzothiazine $Ca^{2+}$ antagonist and related compounds
by A. Ota, S. Ito and Y. Kawashima (Osaka, Japan) . . . . . 37

Simple reversed-phase high-performance liquid chromatographic method for 13-*cis*-retinoic acid in serum
by R. R. Gadde and F. W. Burton (Buffalo, NY, USA) . . . . . 41

Stability studies with a high-performance liquid chromatographic method for the determination of a new anthracycline analogue, 3′-deamino-3′-[2-(*S*)-methoxy-4-morpholino]doxorubicin (FCE 23762), in the final drug formulation
by M. L. Rossini and M. Farina (Milan, Italy) . . . . . 47

Use of direct injection precolumn techniques for the high-performance liquid chromatographic determination of the retinoids acitretin and 13-*cis*-acitretin in plasma
by R. Wyss and F. Bucheli (Basle, Switzerland) . . . . . 55

Determination of the catechol-O-methyltransferase inhibitor Ro 40-7592 in human plasma by high-performance liquid chromatography with coulometric detection
by U. Timm and R. Erdin (Basle, Switzerland) . . . . . 63

High-performance liquid chromatography method for the determination of aminoglycosides based on automated pre-column derivatization with *o*-phthalaldehyde
by M. C. Caturla and E. Cusido (Barcelona, Spain) and D. Westerlund (Uppsala, Sweden) . . . . . 69

Determination of Zy 17617B in plasma by solid-phase extraction and liquid chromatography with automated pre-column exchange
by C. Chollet and M. Salanon (Nyon, Switzerland) . . . . . 73

Determination of (*S*)-(−)-cathinone and its metabolites (*R,S*)-(−)-norephedrine and (*R,R*)-(−)-norpseudoephedrine in urine by high-performance liquid chromatography with photodiode-array detection
by K. Mathys and R. Brenneisen (Berne, Switzerland) . . . . . 79

Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection
by H.-J. Helmlin and R. Brenneisen (Berne, Switzerland) . . . . . 87

High-performance liquid chromatographic determination of α-tocopheryl nicotinate in cosmetic preparations
by A. Baruffini, E. De Lorenzi, C. Gandini, M. Kitsos and G. Massolini (Pavia, Italy) . . . . . 95

*Applications: proteins and their constituents*

Rapid determination of amino acids by high-performance liquid chromatography: release of amino acids by perfused rat liver (*Short Communication*)
by F. Zezza, J. Kerner, M. R. Pascale, R. Giannini and E. A. Martelli (Rome, Italy) . . . . . 99

High-performance liquid chromatography of amino acids, peptides and proteins. CXV. Thermodynamic behaviour of peptides in reversed-phase chromatography
by A. W. Purcell, M. I. Aguilar and M. T. W. Hearn (Clayton, Australia) . . . . . 103

Direct determination of kallikrein by high-performance liquid chromatography
by G. Raspi, A. Lo Moro and M. Spinetti (Pisa, Italy) . . . . . 119

Isolation and characterization of α-glucosidase from *Aspergillus niger*
by K. Břízová, B. Králová, K. Demnerová and I. Vinš (Prague, Czechoslovakia) . . . . . 125

*Applications: other biological applications*

Reversed-phase high-performance liquid chromatography–thermospray mass spectrometry of radiation-induced decomposition products of thymine and thymidine
by M. Berger and J. Cadet (Grenoble, France) and R. Berube, R. Langlois and J. E. van Lier (Québec, Canada) . . . 133

Improved determination of individual molecular species of phosphatidylcholine in biological samples by high-performance liquid chromatography with internal standards
by A. Cantafora and R. Masella (Rome, Italy) . . . . . 139

Determination of benzene metabolites in urine of mice by solid-phase extraction and high-performance liquid chromatography
by H. Schad, F. Schäfer, L. Weber and H. J. Seidel (Ulm, Germany) . . . . . 147

Determination of 5-hydroxymethylfurfural by ion-exclusion chromatography with UV detection (*Short Communication*)
by H.-J. Kim and M. Richardson (Natick, MA, USA) . . . . . 153

Determination of inositol polyphosphates from human T-lymphocyte cell lines by anion-exchange high-performance liquid chromatography and post-column derivatization
by A. H. Guse and F. Emmrich (Erlangen, Germany) . . . . . 157

Separation of flavonol-2-O-glycosides from *Calendula officinalis* and *Sambucus nigra* by high-performance liquid and micellar electrokinetic capillary chromatography
by P. Pietta, A. Bruno and P. Mauri (Milan, Italy) and A. Rava (Pavia, Italy) . . . . . 165

High-performance liquid chromatographic method with fluorescence detection for the determination of total homocyst(e)ine in plasma
by I. Fermo, C. Arcelloni, E. De Vecchi, S. Vigano' and R. Paroni (Milan, Italy) . . . . . 171

High-performance liquid chromatographic analysis of the pigments of blood-red prickly pear (*Opuntia ficus indica*)
by E. Forni, A. Polesello, D. Montefiori and A. Maestrelli (Milan, Italy) . . . . . 177

Determination of catecholamines in urine by liquid chromatography and electrochemical detection after on-line sample purification on immobilized boronic acid
by B.-M. Eriksson and M. Wikström (Mölndal, Sweden) . . . . . 185

Determination of cinnamic acid and paeoniflorin in traditional chinese medicinal preparations by high-performance liquid chromatography
by K.-C. Wen, C.-Y. Huang and F.-S. Liu (Taipei, Taiwan) . . . . . 191

Analysis and isolation of indole alkaloids of fungi by high-performance liquid chromatography
by M. Wurst, R. Kysilka and T. Koza (Prague, Czechoslovakia) . . . . . 201

Use of high-performance liquid chromatographic peak deconvolution and peak labelling to identify antiparasitic components in plant extracts
by N. Perez-Souto (Havana, Cuba) and R. J. Lynch, G. Measures and J. T. Hann (Cambridge, UK) . . . . . 209

Improved high-performance liquid chromatographic method for the determination of coenzyme $Q_{10}$ in plasma
by G. Grossi, A. M. Bargossi, P. L. Fiorella and S. Piazzi (Bologna, Italy), M. Battino (Ancona, Italy) and G. P. Bianchi (Bologna, Italy) . . . . . 217

*Applications: miscellaneous*

Reversed-phase high-performance liquid chromatography of anionic and ethoxylated non-ionic surfactants and pesticides in liquid pesticide formulations
by T. Bán (Budapest, Hungary) and E. Papp and J. Inczédy (Veszprém, Hungary) . . . . . 227

Analysis of block copolymers by high-performance liquid chromatography under critical conditions
by T. M. Zimina, J. J. Kever and E. Yu. Melenevskaya (St. Petersburg, USSR) and A. F. Fell (Bradford, UK) . . . 233

Comparison of several methods for the determination of trace amounts of polar aliphatic monocarboxylic acids by high-performance liquid chromatography
by A. J. J. M. Coenen, M. J. G. Kerkhoff, R. M. Heringa and Sj. van der Wal (Geleen, Netherlands) . . . . . 243

ELECTROPHORESIS AND ELECTROKINETIC CHROMATOGRAPHY

Planar chips technology for miniaturization and integration of separation techniques into monitoring systems. Capillary electrophoresis on a chip
by A. Manz, D. J. Harrison, E. M. J. Verpoorte, J. C. Fettinger, A. Paulus, H. Lüdi and H. M. Widmer (Basle, Switzerland) . . . . . 253

Separation of leucinostatins by capillary zone electrophoresis
by M. G. Quaglia, S. Fanali and A. Nardi (Rome, Italy) and C. Rossi and M. Ricci (Perugia, Italy) . . . . . 259

Separation of naphthalene-2,3-dicarboxaldehyde-labeled amino acids by high-performance capillary electrophoresis with laser-induced fluorescence detection
by T. Ueda, R. Mitchell, F. Kitamura, T. Metcalf and T. Kuwana (Lawrence, KS, USA) and A. Nakamoto (Kyoto, Japan) . . . . . 265

Determination of substituted purines by body fluids by micellar electrokinetic capillary chromatography with direct sample injection
by W. Thormann, A. Minger, S. Molteni, J. Caslavska and P. Gebauer (Bern, Switzerland) . . . . . 275

Factors affecting the separation of inorganic metal cations by capillary electrophoresis
by A. Weston and P. R. Brown (Kingston, RI, USA) and P. Jandik, W. R. Jones and A. L. Heckenberg (Milford, MA, USA) . . . . . 289

Prediction of migration behavior of oligonucleotides in capillary gel electrophoresis
by A. Guttman, R. J. Nelson and N. Cooke (Palo Alto, CA, USA) . . . . . 297

Capillary electrophoresis with electrochemical detection employing an on-column Nafion joint
by T. J. O'Shea, R. D. Greenhagen, S. M. Lunte and C. E. Lunte (Lawrence, KS, USA), M. R. Smyth (Dublin, Ireland), D. M. Radzik (Kansas City, MO, USA) and N. Watanabe (Tokyo, Japan) . . . . . 305

Electrokinetic reversed-phase chromatography with packed capillaries
by H. Yamamoto, J. Baumann and F. Erni (Basle, Switzerland) . . . . . 313

OTHER SEPARATION METHODS

Analytical study of heavy crude oil fractions by coupling of the transalkylation reaction with supercritical fluid chromatography
by P.-L. Desbène and A. Abderrezag (Evreux, France) and B. Desmazières (Paris, France) . . . . . 321

Retention behaviour of closely related coumarins in thin-layer chromatographic preassays for high-performance liquid chromatography according to the "PRISMA" model
by P. Härmälä, H. Vuorela, E.-L. Rahko and R. Hiltunen (Helsinki, Finland) . . . . . 329

Asymmetric-channel flow field-flow fractionation with exponential force-field programming
by J. J. Kirkland, C. H. Dilks, Jr., S. W. Rementer and W. W. Yau (Wilmington, DE, USA) . . . . . 339

Simple solution of velocity profiles of laminar flows in channels of various cross-sections used in field-flow fractionation
by J. Pazourek and J. Chmelík (Brno, Czechoslovakia) . . . . . 357

Separation of pristinamycins by high-speed counter-current chromatography. I. Selection of solvent system and preliminary preparative studies
by S. Drogue, M.-C. Rolet, D. Thiébaut and R. Rosset (Paris, France) . . . . . 363

*Author Index Vols. 592 and 593* . . . . . 373

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*).